

Type III and II universal spacetimes

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2015 J. Phys.: Conf. Ser. 600 012066

(<http://iopscience.iop.org/1742-6596/600/1/012066>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 152.94.27.12

This content was downloaded on 07/01/2016 at 10:05

Please note that [terms and conditions apply](#).

Type III and II universal spacetimes

S. Hervik¹, V. Pravda², A. Pravdová²

¹ Faculty of Science and Technology, University of Stavanger, N-4036 Stavanger, Norway;

² Institute of Mathematics, Academy of Sciences of the Czech Republic
Žitná 25, 115 67 Prague 1, Czech Republic

E-mail: sigbjorn.hervik@uis.no, pravda@math.cas.cz, pravdova@math.cas.cz

Abstract. We briefly summarize our recent results on universal spacetimes. We show that universal spacetimes are necessarily CSI, i.e. for these spacetimes, all curvature invariants constructed from the Riemann tensor and its derivatives are constant. Then, we focus on type III universal spacetimes and discuss a proof of universality for a class of type III Kundt spacetimes. We also mention explicit examples of type III and II universal spacetimes.

1. Introduction

In the contribution [1] in this volume, we have introduced universal spacetimes obeying the following definition [2]

Definition 1.1. A metric is called *universal* if all conserved symmetric rank-2 tensors constructed from the metric, the Riemann tensor and its covariant derivatives of arbitrary order are multiples of the metric.

We have argued that universal metrics solve the vacuum equations of *all* theories of gravity with the Lagrangian of the form

$$L = L(g_{ab}, R_{abcd}, \nabla_{a_1} R_{bcde}, \dots, \nabla_{a_1 \dots a_p} R_{bcde}). \quad (1)$$

In general, it seems too difficult to study universal spacetimes in arbitrary spacetime dimension in full generality. However, employing the algebraic classification of the Weyl tensor [3] (see also [4] for a recent review) leads, in particular for types N and III, to considerable simplification of the problem and allows us to prove universality for various classes of metrics.

In [1], we summarized the main results of [5] for type N universal spacetimes. Here, let us briefly present further results on universal spacetimes (for details see [5]).

First, in section 2, we describe the main points of the proof of the following theorem.

Theorem 1.2. *A universal spacetime is necessarily a CSI spacetime.*

CSI (constant curvature invariant) spacetimes are spacetimes that have all curvature invariants constructed from the metric, the Riemann tensor and its derivatives of arbitrary order constant, see e.g. [6].

Then, we discuss type III spacetimes. Although we expect that type III universal spacetimes are necessarily Kundt, in contrast to the type N case, we cannot use Theorem 1.2 to prove this statement in full generality and in [5], we provide a proof only in the “generic” case. Thus, in



section 3, we focus on the sufficient conditions for universality. We identify a universal subclass of type III Kundt metrics¹

Theorem 1.3. *Type III, $\tau_i = 0$ Einstein Kundt spacetimes obeying*

$$C_{acde}C_b{}^{cde} = 0 \quad (2)$$

are universal.

Finally, in section 4, we present explicit examples of type III and type II universal metrics.

2. Universal spacetimes are CSI

Let us here briefly summarize the main points of the proof of the theorem 1.2.

Let us consider the Lagrangian (1) containing covariant derivatives of the Riemann tensor up to a fixed p . In [7], it has been shown that by varying the action, one arrives at the field equations

$$\begin{aligned} -T^{ab} &= \frac{\partial L}{\partial g_{ab}} + E^a{}_{cde}R^{bcde} + 2\nabla_c\nabla_d E^{acdb} + \frac{1}{2}g^{ab}L, \\ E^{bcde} &= \frac{\partial L}{\partial R_{bcde}} - \nabla_{a_1} \frac{\partial L}{\partial \nabla_{a_1} R_{bcde}} + \dots + (-1)^p \nabla_{(a_1} \dots \nabla_{a_p)} \frac{\partial L}{\partial \nabla_{(a_1} \dots \nabla_{a_p)} R_{bcde}}, \end{aligned} \quad (3)$$

where T^{ab} is the associated conserved tensor.

Now, let us consider any polynomial invariant I and let us assume that it contains derivatives of the Riemann tensor of orders at most p . By [7], we can assume it is of the form

$$I = I[g_{ab}, R_{abcd}, \nabla_{a_1} R_{bcde}, \dots, \nabla_{(a_1 \dots a_p)} R_{bcde}].$$

Let us consider the (infinite) series of Lagrangians $L = I^n$, $n = 1, 2, 3, \dots$. By variation, we get a conserved tensor $T[n]^a{}_b$ for each n . For universal spacetimes, traces of $T[n]^a{}_b$ are constant. By studying the explicit forms of the corresponding expressions [5], one arrives at a conclusion that I has to be a constant and since I is an arbitrary curvature invariant it follows that universal spacetimes are CSI.

3. Type III universal spacetimes

Type III spacetimes by definition [3,4] admit a multiple Weyl aligned null direction, mWAND, ℓ . Let us complete a frame with another null vector n and $n - 2$ spacelike orthonormal vectors $m^{(i)}$ with the only non-vanishing products being $\ell^a n_a = 1$, $m^{(i)a} m_a^{(j)} = \delta_{ij}$ (coordinate indices $a, b, \dots = 0 \dots n - 1$, frame indices $i, j, \dots = 2 \dots n - 1$).

In an appropriately chosen frame, the Weyl tensor for type III spacetime reads [3,4]

$$C_{abcd} = 8\Psi'_i \ell_{\{a} n_b \ell_c m_d^{(i)}\} + 4\Psi'_{ijk} \ell_{\{a} m_b^{(i)} m_c^{(j)} m_d^{(k)}\} + 4\Omega'_{ij} \ell_{\{a} m_b^{(i)} \ell_c m_d^{(j)}\}, \quad (4)$$

where the frame components satisfy $\Psi'_i = \Psi'_{jij}$, $\Omega'_{ij} = \Omega'_{ji}$, and $\Omega'_{ii} = 0$ and for an arbitrary tensor T_{abcd}

$$T_{\{abcd\}} \equiv \frac{1}{2}(T_{[ab][cd]} + T_{[cd][ab]}). \quad (5)$$

As discussed in [5], for type III Einstein spacetimes, the symmetric rank-2 tensor

$$S_{ab} \equiv C_{acde}C_b{}^{cde} \quad (6)$$

is conserved. While for type N, S_{ab} vanishes identically, for type III Einstein spacetimes, it is in general a boost weight -2 tensor proportional to $\ell_a \ell_b$. Therefore, for type III universal spacetimes, we have an additional necessary condition (2).

¹ However, note that other type III Einstein Kundt spacetimes may also exist.

3.1. Main points of the proof of the theorem 1.3

The key intermediate result proven in [5] is that

Proposition 3.1. *For type III Einstein Kundt spacetimes, the boost order of $\nabla^{(k)}C$ (a covariant derivative of an arbitrary order of the Weyl tensor) with respect to the multiple WAND is at most -1 .*

Proof of this result relies on the precise form of various Bianchi and Ricci identities [8,9].

A direct consequence of proposition 3.1 is that

Lemma 3.2. *For type III Ricci-flat Kundt spacetimes, a non-vanishing rank-2 tensor constructed from the metric, the Weyl tensor and its covariant derivatives of arbitrary order is at most quadratic in the Weyl tensor and its covariant derivatives.*

We find that for type III Ricci-flat Kundt spacetimes, for which the FKWC basis [10] of rank-2, order-6 Weyl polynomials consists of

$$F_1 \equiv C^{pqrs}C_{pqrs;ab}, \quad F_2 \equiv C^{pqrs}{}_{;a}C_{pqrs;b}, \quad F_3 \equiv C^{pqrs}{}_{a;s}C_{pqrb}{}^{;s}, \quad (7)$$

F_1 and F_2 are in general non-vanishing. In our case, F_1 and F_2 are conserved and thus, in general, type III Ricci-flat Kundt spacetimes are not universal. However, both F_1 and F_2 vanish for $\tau_i = 0$. In this case, a rather technical proof [5] allows us to arrive at the theorem 1.3.

4. Kundt spacetimes

Since all universal spacetimes we have discussed belong to the Kundt class, let us briefly discuss Kundt metrics. Kundt spacetimes are spacetimes admitting a null geodetic vector field ℓ with vanishing shear, expansion and twist. In appropriately chosen coordinates,

$$\ell_{a;b} = L_{11}\ell_a\ell_b + \tau_i(\ell_a m_b^{(i)} + m_a^{(i)}\ell_b) \quad (8)$$

and the metric reads [6,11]

$$ds^2 = 2du [dr + H(u, r, x^\gamma)du + W_\alpha(u, r, x^\gamma)dx^\alpha] + g_{\alpha\beta}(u, x^\gamma)dx^\alpha dx^\beta, \quad (9)$$

with $\alpha, \beta, \gamma = 2 \dots n - 1$.

Since universality implies CSI (theorem 1.2) we restrict ourselves to the *Kundt CSI metrics*, where [6,12]

$$\begin{aligned} W_\alpha(u, r, x^\gamma) &= rW_\alpha^{(1)}(u, x^\gamma) + W_\alpha^{(0)}(u, x^\gamma), \\ H(u, r, x^\gamma) &= \frac{r^2}{8} \left(a + W_\alpha^{(1)}W^{(1)\alpha} \right) + rH^{(1)}(u, x^\gamma) + H^{(0)}(u, x^\gamma), \end{aligned} \quad (10)$$

$g_{\alpha\beta}(x^\gamma)$ (note that $g_{\alpha\beta,u} = 0$) is the metric of a locally homogeneous space and a is a constant. Note that (10) are necessary but not sufficient conditions for Kundt CSI.

4.1. type III

Note also that for type III and $\tau_i = 0$, the Bianchi identities imply $\Lambda = 0$ and thus type III, $\tau_i = 0$, Kundt universal spacetimes are in fact Ricci-flat and VSI (vanishing scalar invariants) with a metric of the form [13,14]

$$ds^2 = 2du [dr + H(u, r, x^\gamma)du + W_\alpha(u, r, x^\gamma)dx^\alpha] + \delta_{\alpha\beta}dx^\alpha dx^\beta, \quad (11)$$

with

$$W_2 = 0, \quad (12)$$

$$W_M(u, r, x^\gamma) = W_M^{(0)}(u, x^\gamma), \quad (13)$$

$$H(u, r, x^\gamma) = rH^{(1)}(u, x^\gamma) + H^{(0)}(u, x^\gamma). \quad (14)$$

Since further constraints on H and W_M follow from the Einstein equations we conclude with an explicit example of a four-dimensional type III, $\tau_i = 0$ Ricci-flat Kundt universal metric (expressed in other coordinates) [15]

$$ds^2 = 2dudv - x(v + e^x)du^2 + e^x(dx^2 + e^{-2u}dy^2). \quad (15)$$

4.2. type II

An example of a four-dimensional type II universal Kundt spacetime

$$ds^2 = 2dudv + (-v^2\lambda + H(u, x, y))du^2 + \frac{1}{\lambda}(dx^2 + \sinh^2 x dy^2), \quad \square H = 0 \quad (16)$$

has been given already in [2]. As we will discuss elsewhere, for some type II classes of Kundt spacetimes, universality depends on dimension of the spacetime.

Acknowledgments

The authors acknowledge support from research plan RVO: 67985840 and research grant GAČR 13-10042S.

References

- [1] S. Hervik, V. Pravda, and A. Pravdová. Type N universal spacetimes. Proceedings of the 2014 Spanish Relativity Meeting, to appear in Journal of Physics: Conference Series
- [2] A.A. Coley, G.W. Gibbons, S. Hervik, and C.N. Pope. Metrics with vanishing quantum corrections. *Class. Quantum Grav.*, 25:145017, 2008.
- [3] A. Coley, R. Milson, V. Pravda, and A. Pravdová. Classification of the Weyl tensor in higher dimensions. *Class. Quantum Grav.*, 21:L35–L41, 2004.
- [4] M. Ortaggio, V. Pravda, and A. Pravdová. Algebraic classification of higher dimensional spacetimes based on null alignment. *Class. Quantum Grav.*, 30:013001, 2013.
- [5] S. Hervik, V. Pravda and A. Pravdová, Type III and N universal spacetimes. *Class. Quantum Grav.*, 31:215005, 2014.
- [6] A. Coley, S. Hervik, and N. Pelavas. On spacetimes with constant scalar invariants. *Class. Quantum Grav.*, 23:3053–3074, 2006.
- [7] V. Iyer and R. M. Wald, Some Properties of Noether Charge and a Proposal for Dynamical Black Hole Entropy *Phys. Rev. D*, 50: 846-864 , 1994.
- [8] M. Ortaggio, V. Pravda, and A. Pravdová. Ricci identities in higher dimensions. *Class. Quantum Grav.*, 24:1657–1664, 2007.
- [9] M. Durkee, V. Pravda, A. Pravdová, and H. S. Reall. Generalization of the Geroch-Held-Penrose formalism to higher dimensions. *Class. Quantum Grav.*, 27:215010, 2010.
- [10] S. A. Fulling, R. C. King, B.G. Wybourne, and C.J. Cummins. Normal forms for tensor polynomials. 1: The Riemann tensor. *Class. Quantum Grav.*, 9:1151–1197, 1992.
- [11] A. Coley, R. Milson, N. Pelavas, V. Pravda, A. Pravdová, and R. Zalaletdinov. Generalizations of pp -wave spacetimes in higher dimensions. *Phys. Rev. D*, 67:104020, 2003.
- [12] A. Coley, S. Hervik, G. O. Papadopoulos, and N. Pelavas. Kundt spacetimes. *Class. Quantum Grav.*, 26:105016, 2009.
- [13] A. Coley, R. Milson, V. Pravda, and A. Pravdová. Vanishing scalar invariant spacetimes in higher dimensions. *Class. Quantum Grav.*, 21:5519–5542, 2004.
- [14] A. Coley, A. Fuster, S. Hervik, and N. Pelavas. Higher dimensional VSI spacetimes. *Class. Quantum Grav.*, 23:7431–7444, 2006.
- [15] A. Z. Petrov. Gravitational field geometry as the geometry of automorphisms. In *Recent Developments in General Relativity*, pages 379–386. Pergamon Press/PWN, Oxford/Warszawa, 1962.