

# The effect of Zhang's guideline versus the WHO partograph on childbirth experience measured by the Childbirth Experience Questionnaire in the Labor Progression Study (LaPS): A cluster randomized trial

Daniella Judit Rozsa<sup>1,2</sup>  | Rebecka Dalbye<sup>3,4</sup>  | Stine Bernitz<sup>3,4</sup> | Ellen Blix<sup>4</sup> | Ingvild Dalen<sup>5</sup> | Geir Sverre Braut<sup>5</sup>  | Torbjørn M. Eggebø<sup>1,6</sup> | Pål Øian<sup>7</sup> | Ragnar Kvie Sande<sup>1,8</sup>

<sup>1</sup>Department of Obstetrics and Gynecology, Stavanger University Hospital, Stavanger, Norway

<sup>2</sup>Faculty of Health Sciences, Department of Caring and Ethics, University of Stavanger, Stavanger, Norway

<sup>3</sup>Department of Obstetrics and Gynecology, Østfold Hospital Trust, Grålum, Norway

<sup>4</sup>Department of Nursing and Health Promotion, Faculty of Health Sciences, OsloMet - Oslo Metropolitan University, Oslo, Norway

<sup>5</sup>Department of Research, Stavanger University Hospital, Stavanger, Norway

<sup>6</sup>National Center for Fetal Medicine, St. Olav's University Hospital, Trondheim, Norway

<sup>7</sup>Department of Obstetrics and Gynecology, University Hospital of North Norway, Tromsø, Norway

<sup>8</sup>Department of Clinical Science, University of Bergen, Bergen, Norway

## Correspondence

Daniella Judit Rozsa, Department of Obstetrics and Gynecology, Stavanger University Hospital, Helse Stavanger HF, PO Box 8100, 4068 Stavanger, Norway. Email: rodani@sus.no

## Funding information

Sykehuset Østfold Hospital Trust (Grant AB3293), University of Stavanger (Grant HV-AOE30087843).

## Abstract

**Introduction:** Childbirth experience is an increasingly recognized and important measure of quality of obstetric care. Previous research has shown that it can be affected by intrapartum care and how labor is followed. A partograph is recommended to follow labor progression by recording cervical dilation over time. There are currently different guidelines in use worldwide to follow labor progression. The two main ones are the partograph recommended by the World Health Organization (WHO) based on the work of Friedman and Philpott and a guideline based on Zhang's research. In our study we assessed the effect of adhering to Zhang's guideline or the WHO partograph on childbirth experience. Zhang's guideline describes expected normal labor progression based on data from contemporary obstetric populations, resulting in an exponential progression curve, compared with the linear WHO partograph. The choice of labor curve affects the intrapartum follow-up of women and this could potentially affect childbirth experience.

**Material and methods:** The Labor Progression Study (LaPS) study was a prospective, cluster randomized controlled trial conducted at 14 birth centers in Norway. Birth centers were randomized to either follow Zhang's guideline or the WHO partograph. Nulliparous women in active labor, with one fetus in cephalic presentation at term and spontaneous labor onset were included. At 4 weeks postpartum, included women received an online login to complete the Childbirth Experience Questionnaire (CEQ). Total score on the CEQ, the four domain scores on the CEQ, and scores on the individual items on the CEQ were compared between the two groups.

**Abbreviations:** BMI, body mass index; CEQ, Childbirth Experience Questionnaire; CI, confidence interval; LaPS, Labor Progression Study; OR, odds ratio; RM, ratios of means; WHO, World Health Organization.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2021 The Authors. *Acta Obstetrica et Gynecologica Scandinavica* published by John Wiley & Sons Ltd on behalf of Nordic Federation of Societies of Obstetrics and Gynecology (NFOG).

**Results:** There were 1855 women in the Zhang group and 1749 women in the WHO partograph group. There was no difference in the total or domain CEQ scores between the two groups. We found statistically significant differences for two individual items; women in the Zhang group scored lower on positive memories and feeling of control.

**Conclusions:** Based on our findings on childbirth experience there is no reason to prefer Zhang's guideline over the WHO partograph.

**KEYWORDS**

childbirth experience, labor progression, obstetric, WHO partograph, Zhang's guideline

## 1 | INTRODUCTION

Intrapartum care for a positive childbirth experience, published in 2018 by the World Health Organization (WHO), emphasizes the importance of a positive childbirth experience and has identified important knowledge gaps regarding progress of labor and care throughout labor and birth.<sup>1</sup> Childbirth experience may have immediate as well as long-term positive or negative effects on well-being and health. A positive childbirth experience might contribute to an increase in vaginal deliveries, whereas negative experience can contribute to future fear of giving birth. A systematic review confirms the impact of a negative childbirth experience on subsequent reproductive decisions. It is associated with not having another child, delaying the birth of a subsequent child and a preference for cesarean section in subsequent pregnancies.<sup>2,3</sup> One of the factors that influences childbirth, and probably childbirth experience, is how labor progression is monitored.<sup>4</sup> Friedman pioneered research in labor progression by graphically describing the cervical dilation over time.<sup>5</sup> Based on this concept, Philpott developed guidelines for graphically assessing labor progression constructing the partograph in 1972.<sup>6,7</sup> In 1994, their work became the basis of the WHO partograph that have been used worldwide since.<sup>8</sup> At the start of active labor, a timeline is placed on the woman's partograph. The linear curve of expected labor progression is constant throughout labor and serves as a reference point for labor dystocia. Due to changes in clinical practices and obstetric populations during the past decades, the use of the WHO partograph in contemporary obstetric populations has been questioned.<sup>9</sup> The guidelines were thought to overestimate the expected speed of cervical dilation compared with the physiological rate in current obstetric populations, thus leading to increased interventions and intrapartum cesarean section rates.<sup>9</sup> In 2010, Zhang et al. presented a labor curve based on a large contemporary cohort.<sup>10</sup> They found an initial slower labor progression pattern that followed an exponential curve and did not find the previously described deceleration phase. Zhang's approach calls for an evaluation of the woman throughout labor based on her current cervical status, and thus seems to follow labor physiology more than the previously established linear cut-offs. A recent Cochrane review concludes that there is no consensus concerning which partograph is most beneficial or optimal.<sup>11</sup> Recently, the

### Key message

Childbirth experience was compared in women following the WHO partograph or Zhang's guideline for labor progression. Childbirth experience was not significantly different between these groups. Type of partograph used to follow labor progression did not affect childbirth experience.

mathematical methods used by Zhang have been criticized for not being appropriate for describing labor progression data.<sup>12</sup> The WHO requested more studies to fill the knowledge gaps regarding labor progression management. The Labor Progression Study (LaPS) was conducted as the first randomized controlled trial to investigate maternal and neonatal clinical consequences of using a guideline based on Zhang's normal labor curve compared with the WHO partograph that was recommended between 1997 and 2020.<sup>13,14</sup> The Childbirth Experience Questionnaire (CEQ) is developed to assess the multi-dimensional nature of women's childbirth experience.<sup>15</sup> This instrument is thoroughly validated, and easily converted to Norwegian, as it was originally developed in Swedish. An optimal tool for assessing labor dystocia may be an important factor in improving childbirth experience, as both unnecessary intervention and failure to take action in prolonged labor may affect childbirth experience. The present study, the LaPS CEQ, is conducted to assess the effect of adhering to Zhang's guideline vs the WHO partograph on childbirth experience. We hypothesized that the use of Zhang's guideline would improve childbirth experience as it follows the physiological labor curve and evaluates the woman's progression individually throughout labor. Information on this aspect of obstetric care can help guide recommendations in the ongoing discussion on which partograph is the most beneficial to use.

## 2 | MATERIAL AND METHODS

LaPS, a cluster randomized controlled multicenter study, was conducted in 14 birth centers in Norway between 1 December 2014

and 31 January 2017. Centers with more than 500 deliveries per year were eligible for inclusion and were stratified according to number of deliveries and intrapartum cesarean section rate. Based on mathematical calculations, Zhang presented expected time intervals from one integer centimeter to the next in active labor.<sup>10</sup> Women who had a progression slower than the 95th percentile of Zhang's defined normal labor curve were diagnosed with labor dystocia. This labor progression assessment tool is denoted as Zhang's guideline in the LaPS study. The birth centers were randomized to either follow Zhang's guideline or the WHO partograph, as recommended at the time of the study (WHO 1997–2020). The study randomized birth centers with respect to guidelines assessing labor dystocia. Once diagnosed, labor dystocia was treated according to national best practice guidelines. Nulliparous women in active labor were enrolled in the study if they had a single fetus in cephalic presentation at term, had a spontaneous labor start and could understand and read Norwegian. Upon inclusion women consented to supplying their sociodemographic characteristics as well as labor data. Women were asked to enter a valid e-mail address on the consent form, in order to receive an online login ID to answer questions about their birth experience, 4 weeks postpartum. The CEQ version 1 was used to measure childbirth experience.<sup>15</sup> The CEQ was developed and validated in Sweden.<sup>15</sup> The CEQ questionnaire was translated from Swedish to Norwegian according to the good practice for translation and cultural adaptation process for patient-reported outcomes.<sup>16</sup> The CEQ consists of 22 items related to childbirth experience, which are categorized into four domains: own capacity, professional support, perceived safety and participation. The first 19 items are to be answered on a four-point Likert scale from 1 to 4, with category 1 denoting a total agreement and category 4 a total disagreement with the statement. Three items are to be answered on a numerical rating scale with scores 0–100, categorized as 1 (0–40), 2 (41–60), 3 (61–80) and 4 (81–100). The total CEQ score is the mean score of all the items and ranges from 1 to 4. In our analyses, this was rescaled to a scoring range of 0 to 100. Domain scores are calculated as means of included items. We compared the total CEQ scores, as well as scores for all four domains and all of the individual items between the two groups. A web-based version of the CEQ was developed at the Department for Research Computing, USIT University of Oslo, Norway, approved by the developer of this instrument and used in this study. Data were stored at the Services for Sensitive Research Data at the Department for Research Computing, USIT University of Oslo.

The core outcome set of the LaPS study, evaluating the use of Zhang's guideline vs the WHO partograph was recording obstetric and neonatal outcomes, as well as childbirth experience measured by the CEQ questionnaire. Comparisons of obstetric and neonatal outcomes between the two groups have been published separately.<sup>13</sup>

## 2.1 | Statistical analyses

Descriptive statistics are given as means and standard deviations for continuous variables and as counts and percentages for categorical


variables. Total CEQ scores and subscale scores were compared between groups by means of generalized linear models with a log link (ie Poisson regression), from which we present estimated ratios of means with 95% confidence intervals (CI) based on robust (sandwich) standard errors, and with *p*-values from Wald tests. Clustering within hospitals was allowed for by including a random intercept term. Since these outcomes were left-skewed, they were reversed before analysis, so that high scores are interpretable as high dissatisfaction/negative experiences. Individual item scores were compared using ordered logistic regression, from which we present estimated odds ratios with 95% CI and *p*-values from Wald tests, applying cluster-robust standard errors. Descriptive statistics were assessed using SPSS 26.0.0.1 (IBM Corp.). All regression analyses were performed in STATA v. 16.1 with functions *mepoisson* and *ologit*. To allow for multiplicity, only *p*-values <0.01 were considered statistically significant.

## 2.2 | Ethical approval

The study protocol was approved by the Regional Committee for Medical and Health Research Ethics on 11 December 2013 in line with the revised Declaration of Helsinki (2013/1862/REK Sør-Øst) and published.<sup>14</sup> The LaPS Study is registered with the ClinicalTrials.gov, registration number NCT02221427 on 20 August 2014. Initial participant enrolment was on 1 December 2014.

## 3 | RESULTS

A total of 7277 women participated in the LaPS. There was no significant difference in intrapartum cesarean section rate between the two guideline groups. In the LaPS CEQ study, 5810 women were invited to participate, of whom 3652 answered the questionnaire, a 62.9% response rate. We removed 48 cases due to incorrect identification numbers, leaving 3604 cases for statistical analysis (Figure 1). Questionnaires were sent out automatically 4 weeks after inclusion, and 1541 (43.5%) participants answered it the same day. 1411 (39.3%) of participants sent their answers back within a week, and 509 (14.4%) participants within a month. Another 71 answers came in in the next 8 months, for the remaining 72 women this information is not registered. Sociodemographic characteristics of women in the two groups were similar to each other, with the exception of relationship status (Table 1). There were a greater number of single mothers in the Zhang group. We also compared our responder cohort (*n* = 3652) with the non-responder group (*n* = 3625), and found that the sociodemographic characteristics of these two groups did not differ significantly. We have also compared the frequency of intrapartum C-sections, vacuum deliveries and postpartum hemorrhage over 1000 mL between our responder cohort and the non-responder group, and found no significant differences (Tables S1 and S2).



**FIGURE 1** Overview of the inclusion of the Labor Progression Study Childbirth Experience Questionnaire participants [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]

The mean for the CEQ total score and the means for the four domains were comparable between the groups, and we found no statistically significant differences (Table 2). When we compared the odds for higher levels of satisfaction on the individual items of the CEQ, women in the Zhang group scored significantly lower on positive memories and feeling of control (Table 3). Adjustment for relationship status did not change the estimated differences between the groups to any substantial degree for the findings reported in Tables 2 and 3.

## 4 | DISCUSSION

We found no difference in the total and domain CEQ scores for women in the Zhang vs the WHO group. Women allocated to use Zhang's guideline scored lower on positive memories and feeling of control during childbirth than women allocated to use the WHO partograph, and the differences between the two groups were statistically significant. Both of these items are on the perceived safety subscale. Women in our study reported overall good birth experience.

Previous studies show that cesarean section and instrumental delivery rates, as well as medical interventions and adverse neonatal outcomes, all influence childbirth experience. These factors were all primary and secondary outcomes in the LaPS study, and their occurrence did not differ significantly between the two guideline groups.<sup>13,17-19</sup> This can explain the similarities in childbirth

experience we found between the groups. However, we also found differences that could be clinically relevant. An experienced level of control during labor has previously been described as an important factor of determining childbirth experience,<sup>20</sup> and we found this parameter to be significantly better for women in the WHO group compared with the Zhang group. One possible explanation is that Zhang's partograph allows longer time intervals between vaginal examinations before 6 cm of dilation. Thus, women possibly receive less attention and less contact with midwives in this part of labor. There are positive sides to limiting the number of vaginal examinations, yet women may feel less cared for in this scenario, especially if they later experience labor dystocia. Furthermore, following the WHO partograph, women with labor dystocia will cross an alert line prior to crossing the action line. This means intensified follow-up and care at an earlier point in time, as well as earlier warning of possible upcoming labor dystocia. The woman may in this scenario have time to prepare mentally for labor dystocia and its consequences. Women following Zhang's guideline are diagnosed with labor dystocia without previous warning, when reaching a pre-set point in time. This may lead to the labor quite suddenly becoming reclassified as pathological, thus reducing the sense of control. Less sense of control in turn can also lead to less positive memories from the labor and delivery process. Labor duration has also been indicated to influence childbirth experience.<sup>21-23</sup> A Swedish study found that prolonged labor is independently associated with worsening women's childbirth experience. A Danish study found that women who were in labor for less than 12 hours scored higher on all dimensions

**TABLE 1** Sociodemographic characteristics of women in the Labor Progression Study Childbirth Experience Questionnaire compared between those randomized to the Zhang and those randomized to the WHO group

	Zhang	WHO
Age <sup>a</sup>	27.6 (4.5)	27.7 (4.5)
Age groups (years)		
<25	460 (24.8)	419 (24.0)
25–35	1249 (67.3)	1208 (69.1)
>35	146 (7.9)	122 (7.0)
Relationship status		
Single	112 (6.0)	62 (3.5)
Cohabitant	1159 (62.5)	1269 (72.6)
Married	577 (31.1)	403 (23.0)
Other	7 (0.4)	15 (0.9)
Education		
Elementary school	89 (4.8)	69 (3.9)
High school	635 (34.2)	607 (34.7)
Higher education ≥12 years	1131 (61.0)	1073 (61.3)
Smoking first trimester	106 (5.7)	96 (5.6)
BMI <sup>b</sup>	23.6 (4.3)	23.7 (4.2)
BMI groups		
<18.5	75 (4.0)	74 (4.3)
18.5–24.9	1264 (68.3)	1155 (66.3)
25.0–29.9	359 (19.4)	369 (21.2)
≥30.0	154 (8.3)	143 (8.2)
Gestational age at onset of labor <sup>c</sup>	281.3 (7.7)	281.2 (7.4)

Note: Total  $n = 3604$ . Results presented as  $n$  (%) unless where otherwise indicated.

Abbreviations: BMI, body mass index; WHO, World Health Organization.

<sup>a</sup>Mean maternal age in years (standard deviation).

<sup>b</sup>Mean prepregnancy BMI in  $\text{kg}/\text{m}^2$  (standard deviation).

<sup>c</sup>Mean gestational age at onset of labor in days (standard deviation).

**TABLE 2** Comparison of total and subscale Childbirth Experience Questionnaire (CEQ) scores between the Zhang and the WHO groups

CEQ score	Mean score Zhang	Mean score WHO	RM (95% CI)	$p$
Total	69.5	70.0	1.00 (0.97–1.04)	0.80
Domains				
Own capacity	49.2	50.0	1.01 (0.97–1.04)	0.67
Professional support	89.3	89.3	0.98 (0.89–1.08)	0.69
Perceived safety	75.0	75.6	1.01 (0.95–1.07)	0.84
Participation	78.9	80.0	1.01 (0.96–1.06)	0.81

Note: Results from regression analysis. Absolute scores were converted to a scale 1–100 for comparability. Ratios of means (RM) with 95% confidence intervals (CI). An RM > 1 indicates that the women in the Zhang group were less satisfied than the women in the WHO group. Total  $n = 3604$ : 1749 in the Zhang group, 1855 in the WHO group.

Abbreviations: CEQ, Childbirth Experience Questionnaire; WHO, World Health Organization.

of the CEQ.<sup>24</sup> In the LaPS study, women in the Zhang group had significantly longer overall duration of labor by 0.84 h difference in median. The first stage was longer by 0.66 h, as well as the second stage by 0.18 h in the Zhang group.<sup>25</sup> We cannot rule out an association between duration of labour and childbirth experience in our study.

Our study contributes important information to the ongoing discussion on which partograph is most beneficial to use. Our results support the previous notion that differences that are found in description of the normal labor curve may be merely methodological, and that finding and testing the most optimal partograph should be the primary aim of research.<sup>26,27</sup> The comparison that our study made was between Zhang's guideline and the WHO partograph in clinical use on nulliparous women. Although these partographs are constructed from available cervical dilation and time data differently, the most important difference between these two guidelines is how they define labor dystocia: Zhang's guideline defines pathology at the 95<sup>th</sup> centile, while the WHO partograph defined pathology at the 90<sup>th</sup> centile. It is plausible that the two study groups are actually quite similar, but can have some differences in labor management that affect some aspects of perceived safety in a woman's birth experience.

Strengths of our study include its size, the randomization and the careful patient selection, including only nulliparous women, which reduces the risk of type 1 error and confounding bias. Another strength is that women were asked to report their childbirth experience 4 weeks postpartum: a point in time when the new mothers are settled into their new roles, have a little distance from labor and delivery, but close enough to still have detailed memory of it. In all, 82.8% of our responders sent in their answers within a week. However, this can also be viewed as a weakness of the study. Previous research has shown that childbirth experience of women can change over time, both for the positive and the negative,<sup>28–30</sup> and the optimal point in time for measuring childbirth experience remains unknown. Another weakness of our study is the relatively low response rate. Furthermore, looking at differences between the groups at the level of individual questions on a questionnaire

CEQ item	OR (95% CI)	p
Labor and birth went as I had expected	0.90 (0.78–1.05)	0.17
I felt strong during labor and birth	0.96 (0.79–1.17)	0.72
I felt scared during labor and birth	1.00 (0.92–1.08)	0.90
I felt capable during labor and birth	0.99 (0.88–1.12)	0.92
I was tired during labor and birth	0.96 (0.82–1.13)	0.64
I felt happy during labor and birth	0.94 (0.88–1.01)	0.11
I have many positive memories from childbirth	0.89 (0.82–0.97)	0.005
I have many negative memories from childbirth	0.96 (0.85–1.08)	0.48
Some of my memories from childbirth make me feel depressed	1.02 (0.87–1.20)	0.80
I felt I had a say whether I could be up and about or lie down	0.97 (0.86–1.09)	0.59
I felt I had a say in deciding my birthing position	0.96 (0.91–1.02)	0.20
I felt I had a say in the choice of pain relief	0.98 (0.89–1.07)	0.59
My midwife devoted enough time to me	0.90 (0.78–1.04)	0.14
My midwife devoted enough time to my partner	0.94 (0.80–1.11)	0.49
My midwife kept me informed about what was happening during labor and birth	1.02 (0.90–1.17)	0.74
My midwife understood my needs	0.96 (0.85–1.09)	0.56
I felt very well cared for by my midwife	0.97 (0.81–1.16)	0.73
My impression of the team's medical skills made me feel secure	0.99 (0.79–1.24)	0.93
I felt that I handled the situation well	0.94 (0.84–1.06)	0.34
As a whole, how painful did you feel childbirth was?	1.02 (0.86–1.19)	0.86
As a whole, how much control did you feel you had during childbirth?	0.87 (0.78–0.96)	0.007
As a whole, how secure did you feel during childbirth?	0.95 (0.85–1.06)	0.33

Note: Results from regression analysis. Group differences assessed in ordered logistic regression models with cluster-robust standard errors. We present odds ratios (OR) with 95% confidence intervals (CI). OR > 1 indicates that the Zhang group had higher odds of responses indicating a more positive birth experience compared with the WHO group. Total  $n = 3604$ : 1749 in the Zhang group, 1855 in the WHO group.

Abbreviations: CEQ, Childbirth Experience Questionnaire; WHO, World Health Organization.

amounts to multiple hypothesis testing, with the associated risk for incidental findings. This was addressed by lowering the significance level to 0.01, thus lowering the risk for type 1 error.

## 5 | CONCLUSION

Overall score and the four domains of childbirth experience did not show significant differences between women in the Zhang guideline or the WHO partograph group. Women in the Zhang group scored significantly lower on positive memories and feeling of control.

Based on our findings on childbirth experience there is no reason to prefer Zhang's guideline over the WHO partograph. Childbirth experience is an important perinatal outcome, and further research is needed to identify measures to improve it.

**TABLE 3** Comparison of single item Childbirth Experience Questionnaire (CEQ) scores between women in the Zhang and the WHO groups

## AUTHOR CONTRIBUTIONS

All authors interpreted the data. All authors contributed to the manuscript and approved the final version, and accept responsibility for the paper as published. DR took part in writing the protocol, data collection, data acquisition, data analysis and the writing of the paper. RD took part in the study design, data collection and the writing of the paper. SB initiated the trial, took part in the study design, writing the protocol, data collection and the writing of the paper. EB initiated the trial, took part in the study design, data collection, and the writing of the paper. ID did the statistical analysis and took part in the writing of the paper. GSB gave input to the analysis and took part in the writing of the paper. TME took part in data collection and the writing of the paper. PØ initiated the trial, took part in the study design, data collection and the writing of the paper. RKS took part in writing the protocol, the data analysis and the writing of the paper.

## ORCID

Daniella Judit Rozsa  <https://orcid.org/0000-0002-1585-1735>

Rebecka Dalbye  <https://orcid.org/0000-0002-0522-0526>

Geir Sverre Braut  <https://orcid.org/0000-0002-3337-4792>

## REFERENCES

- WHO. WHO recommendations: intrapartum care for a positive childbirth experience 2018. Accessed March 1, 2020. <http://apps.who.int/iris/bitstream/handle/10665/260178/9789241550215-eng.pdf;jsessionid=F1F5397399F448885328491174AC61C6?sequence=1>
- Shorey S, Yang YY, Ang E. The impact of negative childbirth experience on future reproductive decisions: a quantitative systematic review. *J Adv Nurs*. 2018;74:1236-1244.
- Pang MW, Leung TN, Lau TK, Hang Chung TK. Impact of first childbirth on changes in women's preference for mode of delivery: follow-up of a longitudinal observational study. *Birth*. 2008;35:121-128.
- Waldenström U, Rudman A, Hildingsson I. Intrapartum and postpartum care in Sweden: women's opinions and risk factors for not being satisfied. *Acta Obstet Gynecol Scand*. 2006;85:551-560.
- Friedman E. The graphic analysis of labor. *Am J Obstet Gynecol*. 1954;68:1568-1575.
- Philpott RH, Castle WM. Cervicographs in the management of labour in primigravidae. I. The alert line for detecting abnormal labour. *J Obstet Gynaecol Br Commonw*. 1972;79:592-598.
- Philpott RH, Castle WM. Cervicographs in the management of labour in primigravidae. II. The action line and treatment of abnormal labour. *J Obstet Gynaecol Br Commonw*. 1972;79:599-602.
- World Health Organization. World Health Organization partograph in management of labour. Maternal Health and Safe Motherhood Programme. *Lancet*. 1994;1994:1399-1404.
- Zhang J, Troendle JF, Yancey MK. Reassessing the labor curve in nulliparous women. *Am J Obstet Gynecol*. 2002;187:824-828.
- Zhang J, Landy HJ, Ware Branch D, et al. Contemporary patterns of spontaneous labor with normal neonatal outcomes. *Obstet Gynecol*. 2010;116:1281-1287.
- Lavender T, Cuthbert A, Smyth RM. Effect of partograph use on outcomes for women in spontaneous labour at term and their babies. *Cochrane Database Syst Rev*. 2018;(8):CD005461.
- de Vries BS, McDonald S, Joseph FA, et al. Impact of analysis technique on our understanding of the natural history of labour: a simulation study. *BJOG*. 2021;128:1833-1842.
- Bernitz S, Dalbye R, Zhang J, et al. The frequency of intrapartum caesarean section use with the WHO partograph versus Zhang's guideline in the Labour Progression Study (LaPS): a multicentre, cluster-randomised controlled trial. *Lancet*. 2019;393:340-348.
- Bernitz S, Dalbye R, Oian P, Zhang J, Eggebo TM, Blix E. Study protocol: the Labor Progression Study, LAPS—does the use of a dynamic progression guideline in labor reduce the rate of intrapartum cesarean sections in nulliparous women? A multicenter, cluster randomized trial in Norway. *BMC Pregnancy Childbirth*. 2017;17:370.
- Dencker A, Taft C, Bergqvist L, Lilja H, Berg M. Childbirth experience questionnaire (CEQ): development and evaluation of a multi-dimensional instrument. *BMC Pregnancy Childbirth*. 2010;10:81.
- Wild D, Grove A, Martin M, et al. Principles of good practice for the translation and cultural adaptation process for Patient-Reported Outcomes (PRO) measures: report of the ISPOR task force for translation and cultural adaptation. *Value Health*. 2005;8:94-104.
- Soet JE, Brack GA, Dilorio C. Prevalence and predictors of women's experience of psychological trauma during childbirth. *Birth*. 2003;30:36-46.
- Waldenström U, Borg IM, Olsson B, Skold M, Wall S. The childbirth experience: a study of 295 new mothers. *Birth*. 1996;23:144-153.
- Waldenström U, Hildingsson I, Rubertsson C, Radestad I. A negative birth experience: prevalence and risk factors in a national sample. *Birth*. 2004;31:17-27.
- Goodman P, Mackey MC, Tavakoli AS. Factors related to childbirth satisfaction. *J Adv Nurs*. 2004;46:212-219.
- Lavender T, Alfirevic Z, Walkinshaw S. Partogram action line study: a randomised trial. *Br J Obstet Gynaecol*. 1998;105:976-980.
- Lavender T, Wallymahmed AH, Walkinshaw SA. Managing labor using partograms with different action lines: a prospective study of women's views. *Birth*. 1999;26:89-96.
- Adams SS, Eberhard-Gran M, Eskild A. Fear of childbirth and duration of labour: a study of 2206 women with intended vaginal delivery. *BJOG*. 2012;119:1238-1246.
- Boie S, Lauridsen HH, Glavind J, Smed MK, Uldbjerg N, Bor P. The Childbirth Experience Questionnaire (CEQ)—validation of its use in a Danish-speaking population of new mothers stimulated with oxytocin during labour. *PLoS One*. 2020;15:e0233122.
- Dalbye R, Blix E, Frøslie KF, et al. The Labour Progression Study (LaPS): duration of labour following Zhang's guideline and the WHO partograph—a cluster randomised trial. *Midwifery*. 2020;81:102578.
- Cohen WR, Friedman EA. Perils of the new labor management guidelines. *Am J Obstet Gynecol*. 2015;212:420-427.
- Cohen WR, Friedman EA. Misguided guidelines for managing labor. *Am J Obstet Gynecol*. 2015;212(753):e1-e3.
- Donate-Manzanares M, Rodríguez-Cano T, Gómez-Salgado J, et al. Quality of childbirth care in women undergoing labour: satisfaction with care received and how it changes over time. *J Clin Med*. 2019;8:434.
- Waldenström U. Why do some women change their opinion about childbirth over time? *Birth*. 2004;31:102-107.
- Stadlmayr W, Amsler F, Lemola S, et al. Memory of childbirth in the second year: the long-term effect of a negative birth experience and its modulation by the perceived intranatal relationship with caregivers. *J Psychosom Obstet Gynaecol*. 2006;27:211-224.

## SUPPORTING INFORMATION

Additional supporting information may be found in the online version of the article at the publisher's website.

**How to cite this article:** Rozsa DJ, Dalbye R, Bernitz S, et al. The effect of Zhang's guideline versus the WHO partograph on childbirth experience measured by the Childbirth Experience Questionnaire in the Labor Progression Study (LaPS): A cluster randomized trial. *Acta Obstet Gynecol Scand*. 2021;00:1-7. doi:[10.1111/aogs.14298](https://doi.org/10.1111/aogs.14298)